arXiv:1405.4787v2 [math.CO] 3 Jul 2015

The Sparing Number of Certain Graph Powers

N. K. Sudev

Department of Mathematics
Vidya Academy of Science & Technology
Thalakkottukara, Thrissur - 680501, India.
E-mail: sudevnk@gmail.com

K. P. Chithra

Naduvath Mana, Nandikkara
Thrissur - 680301, India.
E-mail: chithrasudev@gmail.com

K. A. Germina

PG & Research Department of Mathematics
Mary Matha Arts & Science College
*Mananthavady, Waynad-670645, E-mail:*srgerminaka@gmail.com

Abstract

Let $\mathbb{N}_0$ be the set of all non-negative integers and $\mathcal{P}(\mathbb{N}_0)$ be its power set. Then, an integer additive set-indexer (IASI) of a given graph G is an injective function $f : V(G) \to \mathcal{P}(\mathbb{N}_0)$ such that the induced function $f^+ : E(G) \to \mathcal{P}(\mathbb{N}_0)$ defined by $f^+(uv) = f(u) + f(v)$ is also injective. An IASI f is said to be a weak IASI if $|f^+(uv)| = max(|f(u)|, |f(v)|)$ for all $u, v \in V(G)$. A graph which admits a weak IASI may be called a weak IASI graph. The set-indexing number of an element of a graph G, a vertex or an edge, is the cardinality of its set-labels. The sparing number of a graph G is the minimum number of edges with singleton set-labels, required for a graph G to admit a weak IASI. In this paper, we study the admissibility of weak IASI by certain graph powers and their corresponding sparing numbers.

Key words: Graph powers, integer additive set-indexers, weak integer additive set-indexers, mono-indexed elements of a graph, sparing number of a graph.

AMS Subject Classification : 05C78

1 Introduction

For all terms and definitions, not defined specifically in this paper, we refer to [3], [7] and [15]. For different graph classes, we further refer to [2], [4] and [16]. Unless mentioned otherwise, all graphs considered here are simple, finite and have no isolated vertices.

The sumset of two non-empty sets A and B is denoted by $A+B$ and is defined by $A+B=\{a+b : a \in A, b \in B\}$. Using the concept of sumsets of two sets we have the following notion.

Let $\mathbb{N}_0$ denote the set of all non-negative integers. An *integer additive set-indexer* (IASI, in short) of a graph G is defined in [5] as an injective function $f : V(G) \to \mathcal{P}(\mathbb{N}_0)$ such that the induced function $f^+ : E(G) \to \mathcal{P}(\mathbb{N}_0)$ defined by $f^+(uv) = f(u) + f(v)$ is also injective.

The cardinality of the labeling set of an element (vertex or edge) of a graph G is called the *set-indexing number* of that element.

Lemma 1.1. [6] *Let A and B be two non-empty finite sets of non-negative integers. Then,* $\max(|A|, |B|) \leq |A + B| \leq |A|\,|B|$. *Therefore, for an integer additive set-indexer f of a graph G, we have* $max(|f(u)|, |f(v)|) \leq |f^+(uv)| = |f(u) + f(v)| \leq |f(u)||f(v)|$, *where $u, v \in V(G)$.*

Definition 1.2. [6] An IASI f is said to be a *weak IASI* if $|f^+(uv)| = |f(u) + f(v)| = \max(|f(u)|, |f(v)|)$ for all $uv \in E(G)$. A graph which admits a weak IASI may be called a *weak IASI graph.*

A weak IASI f is said to be *weakly k-uniform IASI* if $|f^+(uv)| = k$, for all $u, v \in V(G)$ and for some positive integer k.

Lemma 1.3. [6] *An IASI f define on a graph G is a weak IASI of G if and only if, with respect to f, at least one end vertex of every edge of G has the set-indexing number* 1.

Definition 1.4. [9] An element (a vertex or an edge) of graph which has the set-indexing number 1 is called a *mono-indexed element* of that graph. The *sparing number* of a graph G is defined to be the minimum number of mono-indexed edges required for G to admit a weak IASI and is denoted by $\varphi(G)$.

The following are some major results on the spring number of certain graph classes, which are relevant in our present study.

Theorem 1.5. [9] *An odd cycle C_n contains odd number of mono-indexed edges and an even cycle contains an even number of mono-indexed edges.*

Theorem 1.6. [9] *The sparing number of an odd cycle C_n is* 1 *and that of an even cycle is* 0.

Theorem 1.7. [9] *The sparing number of a bipartite graph is* 0.

Theorem 1.8. [9] *The sparing number of a complete graph K_n is* $\frac{1}{2}(n-1)(n-2)$.

Now, let us recall the definition of graph powers.

Definition 1.9. [3] The r-th power of a simple graph G is the graph G^r whose vertex set is V, two distinct vertices being adjacent in G^r if and only if their distance in G is at most r. The graph G^2 is referred to as the *square* of G, the graph G^3 as the *cube* of G.

The following is an important theorem on graph powers.

Theorem 1.10. [14] *If d is the diameter of a graph G, then G^d is a complete graph.*

Some studies on the sparing numbers of certain graph classes and certain graph structures have been done in [11], [12] and [13]. As a continuation to these studies, in this paper we determine the sparing number of the powers certain graph classes.

2 Sparing Number of Square of Some Graphs

In this section, we estimate the sparing number of the square of certain graph classes. It is to be noted that the weak IASI f which gives the minimum number of mono-indexed edges in a given graph G will not induce a weak IASI for its square graph, since some of the vertices having non-singleton set-labels will also be at a distance 2 in G. Hence, interchanging the set-labels or relabeling certain vertices may be required to obtain a weak IASI for the square graph of a given graph.

First consider a path graph P_n on n vertices. The following theorem provides the sparing number of the square of a path P_n.

Proposition 2.1. *Let G be a path graph on n vertices. Then,*

$$\varphi(G) = \begin{cases} \frac{1}{3}(2n-3) & \text{if } n \equiv 0 \ (mod\ 3) \\ \frac{1}{3}(2n-2) & \text{if } n \equiv 1 \ (mod\ 3) \\ \frac{1}{3}(2n-1) & \text{if } n \equiv 2 \ (mod\ 3) \end{cases}$$

Proof. Let $P_m : v_1v_2v_3 \dots v_n$, where $m = n-1$. In P_m^2, $d(v_1) = d(v_n) = 2$ and $d(v_2) = d(v_{n-1}) = 3$ and $d(v_r) = 4$, where $3 \leq r \leq n-2$. Hence, $|E(P_m^2)| = \frac{1}{2}\sum_{v\in V} d(v) = \frac{1}{2}[2 \times 2 + 2 \times 3 + 4(n-4)] = (2n-3)$. Also, for $1 \leq i \leq n-2$, the vertices $v_i, v_{i+1},$ and v_{i+2} form a triangle in P_m^2. Then, by Theorem 1.10, each of these triangles must have a mono-indexed edge. That is, among any three consecutive vertices $v_i, v_{i+1},$ and v_{i+2} of P_m, two vertices must be mono-indexed. We require an IASI which makes the maximum possible number of vertices that are not mono-indexed. Hence, label v_1 and v_2 by singleton sets and v_3 by a non-singleton set. Since v_4 and v_5 are adjacent to v_3, they can be labeled only by distinct singleton sets that are not used before for labeling. Now, v_6 can be labeled by a non-singleton set that has not already been used. Proceeding like

this the vertices which has the form $v_{3k}, 3k \leq n$ can be labeled by distinct non-singleton sets and all other vertices by singleton sets. Now, we have to consider the following cases.

Case-1: If $n \equiv 0 \ (mod\ 3)$, then $n = 3k$. Therefore, v_n can also be labeled by a non-singleton set. Then the number of vertices that are not mono-indexed is $\frac{n}{3}$. Therefore, the number of edges that are not mono-indexed is $4(\frac{n}{3} - 1) + 2 = \frac{1}{3}(4n-6)$. Therefore, the total number of mono-indexed edges is $(2n-3) - \frac{1}{3}(4n-6) = \frac{1}{3}(2n-3)$.

Case-2: If $n \equiv 1 \ (mod\ 3)$, then $n - 1 = 3k$. Then, v_{n-1} can be labeled by a non-singleton set and v_n can be labeled by a singleton set. Then the number of vertices that are not mono-indexed is $\frac{n-1}{3}$. Therefore, the number of edges that are not mono-indexed is $4(\frac{(n-1)}{3} - 1) + 3 = \frac{1}{3}(4n-7)$. Therefore, the total number of mono-indexed edges is $(2n-3) - \frac{1}{3}(4n-7) = \frac{1}{3}(2n-2)$.

Case-3: If $n \equiv 2 \ (mod\ 3)$, then $n - 2 = 3k$. Then, v_{n-2} can be labeled by a non-singleton set and v_n and v_{n-1} can be labeled by distinct singleton sets. Then the number of vertices that are not mono-indexed is $\frac{n-2}{3}$. Therefore, the number of edges that are not mono-indexed is $4(\frac{(n-2)}{3} = \frac{1}{3}(4n-8)$. Therefore, the total number of mono-indexed edges is $(2n-3) - \frac{1}{3}(4n-8) = \frac{1}{3}(2n-1)$. □

Figure 1 illustrates squares of even and odd paths which admit weak IASIs. Mono-indexed edges of the graphs are represented by dotted lines.



Figure 1: Squares of even and odd paths which admit weak IASI

Next, we shall discuss the sparing number of the square of cycles. We have $C_3^2 = C_3 = K_3$, $C_4^2 = K_4$ and $C_5^2 = K_5$ and hence by Theorem 1.8, their sparing numbers are $1, 3$ and 6 respectively. The following theorem determines the sparing number of the square of a given cycle on n vertices, for $n \geq 5$.

Theorem 2.2. *Let C_n be a cycle on n vertices. Then, the sparing number of the*

square of C_n is given by

$$\varphi(C_n^2) = \begin{cases} \frac{2}{3}n & \text{if } n \equiv 0 \ (mod\ 3) \\ \frac{2}{3}(n+2) & \text{if } n \equiv 1 \ (mod\ 3) \\ \frac{2}{3}(n+4) & \text{if } n \equiv 2 \ (mod\ 3). \end{cases}$$

Proof. Let $C_n : v_1v_2v_3 \ldots v_nv_1$ be the given cycle on n vertices. The square of C_n is a 4-regular graph. Also, $V(C_n^2) = V(C_n)$. Therefore, by the first theorem on graph theory, we have $\sum_{v \in V} d(v) = 2|E|$. That is, $2|E| = 4n \Rightarrow |E| = 2n, n \geq 5$.

First, label the vertex v_1 in C_n^2 by a non-singleton set. Therefore, four vertices $v_2, v_3, v_n, \text{and} v_{n-1}$ must be labeled by distinct singleton sets. Next, we can label the vertex v_4 by a non-singleton set, that is not already used for labeling. The vertices v_2 and v_3 have already been mono-indexed and the vertices v_5 and v_6 that are adjacent to v_4 in C_n^2 must be labeled by distinct singleton sets that are not used before for labeling. Proceeding like this, we can label all the vertices of the form v_{3k+1}, where k is a positive integer such that $3k + 1 \leq n - 2$ (since the last vertex that remains unlabeled is v_{n-2}).

Here, we need to consider the following cases.

Case-1: If $n \equiv 0 \ (mod\ 3)$, then $n - 2 = 3k + 1$ for some positive integer k. Then, v_{n-2} can be labeled by a non-singleton set. Therefore, the number of vertices that are labeled by non-singleton sets is $\frac{n}{3}$. Since C_n^2 is 4-regular, we have the number of edges that are not mono-indexed in C_n^2 is $\frac{4n}{3}$. Hence, the number of mono-indexed edges is $2n - \frac{4n}{3} = \frac{2n}{3}$.

Case-2: If $n \equiv 1 \ (mod\ 3)$, then $n - 2 \neq 3k + 1$ for some positive integer k. Then, v_{n-2} can not be labeled by a non-singleton set. Here $n - 3 = 3k + 1$ for some positive integer k. Therefore, the number of vertices that are labeled by non-singleton sets is $\frac{n-1}{3}$ and the number of edges that are not mono-indexed in C_n^2 is $\frac{4(n-1)}{3}$. Hence, the number of mono-indexed edges is $2n - \frac{4(n-1)}{3} = \frac{2(n+2)}{3}$.

Case-3: If $n \equiv 2 \ (mod\ 3)$, then neither $n - 2$ nor $n - 3$ is equal to $3k + 1$ for some positive integer k. Here $n - 4 = 3k + 1$ for some positive integer k. Therefore, the number of vertices that are labeled by non-singleton sets is $\frac{n-2}{3}$ and the number of edges that are not mono-indexed in C_n^2 is $\frac{4(n-2)}{3}$. Hence, the number of mono-indexed edges is $2n - \frac{4(n-2)}{3} = \frac{2(n+4)}{3}$. □

Figure 2 illustrates the admissibility of weak IASIs by the squares of cycles. The graphs given in the figure are examples to the weak IASIs of an even cycle and an odd cycle respectively.

A question that arouses much interest in this context is about the sparing number of the powers of bipartite graphs. Invoking Theorem 1.10, we first verify the existence of weak IASIs for the complete bipartite graphs.

Theorem 2.3. *The sparing number of the square of a complete bipartite graph $K_{m,n}$ is $\frac{1}{2}(m + n - 1)(m + n - 1)$.*

Proof. The diameter of a graph $K_{m,n}$ is 2. Hence by Theorem 1.10, $K_{m,n}^2 = K_{m+n}$. Hence, every pair of vertices, that are not mono-indexed, are at a distance 2. The



Figure 2: Weak IASIs of C_{12}^2 and C_7^2.

set-labels of all these vertices, except one, must be replaced by distinct singleton sets. Therefore, by Theorem 1.8, $\varphi(K_{m,n}^2) = \frac{1}{2}(m+n-1)(m+n-2)$. □

A *balanced bipartite graph* is the bipartite graph which has equal number of vertices in each of its bipartitions.

Corollary 2.4. *If G is a balanced complete bipartite graph on $2n$ vertices, then $\varphi(G) = (n-1)(2n-1)$*

Proof. Let $G = K_{n,n}$. Then by Theorem 2.3, $\varphi(G) = \frac{1}{2}(2n-1)(2n-2) = (n-1)(2n-1)$. □

Let G be a bipartite graph. The vertices which are at a distance 2 are either simultaneously mono-indexed or simultaneously labeled by non-singleton sets. Therefore, in G^2, among any pair of vertices which are are not mono-indexed and are at a distance 2 between them, one vertex should be relabeled by a singleton set. Hence, the sparing number of the square of a bipartite graph G depends on the adjacency pattern of its vertices. Hence, the problem of finding the sparing number of bipartite graphs does not offer much scope in this context.

Now we proceed to study the admissibility of weak IASI by the squares of certain other graph classes. First, we discuss about the sparing number of *wheel graphs*. A wheel graph can be defined as follows.

Definition 2.5. [4] A *wheel graph* is a graph defined by $W_{n+1} = C_n + K_1$. The following theorem discusses the sparing number of the square of a wheel graph W_{n+1}.

The sparing number of the square of a wheel graph W_{n+1} is determined in the following result.

Proposition 2.6. *The sparing number of the square of a wheel graph on $n+1$ vertices is $\frac{1}{2}n(n-1)$.*

Proof. The diameter of a wheel graph W_{n+1}, for any positive integer $n \geq 3$, is 2. Hence, by Theorem 1.10, the square of a wheel graph W_{n+1} is a complete graph on $n+1$ vertices. Therefore, by Theorem 1.8, the sparing number of the square graph W_{n+1}^2 is $\frac{1}{2}n(n-1)$. □

Next, we determine the sparing number of another graph class known as *helm graphs* which is defined as follows.

Definition 2.7. *A* helm graph*, denoted by H_n, is the graph obtained by adjoining a pendant edge to each vertex of the outer cycle C_n of a wheel graph W_{n+1}. It has $2n+1$ vertices and $3n$ edges.*

The following result determines the sparing number of a helm graph.

Theorem 2.8. *The sparing number of the square of a helm graph H_n is $\frac{1}{2}n(n+1)$.*

Proof. Let v be the central vertex, $V = \{v_1, v_2, v_3, \ldots, v_n\}$ be the vertex set of the outer cycle of the corresponding wheel graph and $W = \{w_1, w_2, w_3, \ldots, w_n\}$ be the set of pendant vertices in H_n.

The vertex v is adjacent to all the vertices in V and is at distance 2 from all the vertices in W. Therefore, the degree of v in H_n^2 is $2n$. In H_n, for $1 \leq i \leq n$, each v_i is adjacent to two vertices v_{i-1} and v_{i+1} in V and is adjacent to w_i in W and to the vertex v and is at a distance 2 from all the remaining vertices in V and from the vertices w_{i-1} and w_{i+2} in W. Therefore, the degree of each $v_i \in V$ in H_n^2 is $n+3$. Now, in H_n, each vertex w_i is adjacent to the vertex v_i in V and is at a distance 2 from two vertices v_{i-1} and v_{i+2} in V and to the central vertex v. Hence, the degree of each $w_i \in W$ in H_n^2 is 4. Therefore, the number of edges in H_n, $|E| = \frac{1}{2}\sum_{u \in V(H_n)} d(u) = \frac{1}{2}[2n + n(n+3) + 4n] = \frac{1}{2}n(n+9)$.

It is to be noted that W is an independent set in H_n^2 and we can label all vertices in W by distinct non-singleton sets. It can be seen that there are more edges in H_n^2 that are not mono-indexed if we label all the vertices of W by non-singleton sets than labeling possible number of vertices of $V \cup \{v\}$ by non-singleton sets. Therefore, the number of edges of H_n^2 which are not mono-indexed is $4n$. Therefore, the number of mono-indexed edges in H_n^2 is $\frac{1}{2}n(n+9) - 4n = \frac{1}{2}n(n+1)$. □

Figure 3 illustrates the existence of a weak IASI for the square of a helm graph.

An interesting question in this context is about the sparing number of some graph classes containing complete graphs as subgraphs. An important graph class of this kind is a complete *n-sun* which is defined as follows.

Definition 2.9. [2] An *n-sun* or a *trampoline*, denoted by S_n, is a chordal graph on $2n$ vertices, where $n \geq 3$, whose vertex set can be partitioned into two sets $U = \{u_1, u_2, c_3, \ldots, u_n\}$ and $W = \{w_1, w_2, w_3, \ldots, w_n\}$ such that W is an independent set of G and w_j is adjacent to u_i if and only if $j = i$ or $j = i+1$ $(mod\ n)$. *A complete sun* is a sun G where the induced subgraph $\langle U \rangle$ is complete.

The following theorem determines the sparing number of the square of complete sun graphs.



Figure 3: Square of a helm graph with a weak IASI defined on it.

Theorem 2.10. *Let G be the complete sun graph on $2n$ vertices. Then sparing number of G^2 is*

$$\varphi(G^2) = \begin{cases} n^2+1 & \text{if } n \text{ is odd} \\ \frac{n}{2}(2n-1) & \text{if } n \text{ is even.} \end{cases}$$

Proof. Let G be a sun graph on $2n$ vertices, whose vertex set can be partitioned into two sets $U = \{u_1, u_2, c_3, \ldots, u_n\}$ and $W = \{w_1, w_2, w_3, \ldots, w_n\}$ such that w_j is adjacent to u_i if and only if $j = i$ or $j = i+1$ $(mod\ n)$, where W is an independent set and the induced subgraph $\langle U \rangle$ is complete.

In G, the degree of each u_i is $n+1$ and the degree of each w_j is 2. It can be seen that each vertex w_j is adjacent to two vertices in U and is at a distance 2 from all other vertices in U. Hence, in G^2, each vertex w_j is adjacent to all vertices in U and to two vertices w_{j-1} and w_{j+1} (in the sense that $w_0 = w_n$ and $w_{n+1} = w_1$). That is, in G^2, the degree of each vertex w_j in W is $n+2$ and the degree of each vertex u_i in U is $2n-1$. Therefore, $|E(G^2)| = \frac{1}{2}\sum_{v \in V} d(v) = \frac{1}{2}[n(n+2) + n(2n-1)] = \frac{1}{2}n(3n+1)$.

If we label any vertex u_i by a non-singleton set, then no other vertex in G^2 can be labeled by non-singleton sets, as each u_i is adjacent to all other vertices in G^2. Therefore, we label possible number of vertices in W by non-singleton sets. Since w_j is adjacent to w_{j+1}, only alternate vertices in W can be labeled by non-singleton sets.

Case 1: If n odd, then $\frac{1}{2}(n-1)$ vertices W can be labeled by distinct non-singleton sets. Therefore, the number of edges that are not mono-indexed in G^2 is $\frac{1}{2}(n-1)(n+2)$. Hence, the number of mono-indexed edges in G^2 is $\frac{1}{2}n(3n+1) - \frac{1}{2}(n-1)(n+2) = n^2+1$.

Case 2: If n even, then $\frac{n}{2}$ vertices W can be labeled by distinct non-singleton sets. Therefore, the number of edges that are not mono-indexed in G^2 is $\frac{1}{2}n(n+2)$.

Hence, the number of mono-indexed edges in G^2 is $\frac{1}{2}n(3n+1) - \frac{1}{2}n(n+2) = \frac{1}{2}n(2n-1)$. □

Theorem 2.10 is illustrated in Figure 4. The first and second graphs in 2.10 are example to the weak IASIs of the square of the complete n-sun graphs where n is odd and even respectively.



Figure 4: Weak IASIs of the square of a complete 3-sun and a complete 4-sun.

Another important graph that contains a complete graph as one of its subgraph is a split graph, which is defined as follows.

Definition 2.11. [2] A *split graph* is a graph in which the vertices can be partitioned into a clique K_r and an independent set S. A split graph is said to be a *complete split graph* if every vertex of the independent set S is adjacent to every vertex of the the clique K_r and is denoted by $K_S(r, s)$, where r and s are the orders of K_r and S respectively.

The following theorem establishes the sparing number of the square of a complete split graph.

Theorem 2.12. *Let $G = K_S(r, s)$ be a complete split graph without isolated vertices. Then, the sparing number of G^2 is $\frac{1}{2}[(r+s-1)(r+s-2)]$, where $r = |V(K_r)|$ and $s = |S|$.*

Proof. Since G has no isolated vertices, every vertex of v_i S is adjacent to at least one vertex u_j of K_r. Then, v_i is at a distance 2 from all other vertices of K_r. Hence, in G^2 each vertex v_i in S is adjacent to all the vertices of K_r. Also, in G, two vertices of S is at a distance 2 from all other vertices of S. Therefore, every pair of vertices in S are also adjacent in G^2. That is, G^2 is a complete graph on $r + s$ vertices. Hence, by Theorem 1.8, $\varphi(G^2) = \frac{1}{2}[(r + s - 1)(r + s - 2)$. □

So far we have discussed about the sparing number of square of certain graph classes. In this context, a study about the sparing number of the higher powers of these graph classes is noteworthy. In the following section, we discuss about the sparing number of arbitrary powers of certain graph classes.

3 Sparing Number of Arbitrary Graph Powers

For any positive integer n, we know that the diameter of a complete graph K_n is 1. Hence, any power of K_n, denoted by K_n^r is K_n itself. Hence we have the following result.

Proposition 3.1. *For a positive integer r, $\varphi(K_n^r) = \frac{1}{2}(n-1)(n-2)$.*

Proof. We have $K_n^r = K_n$. Hence, $\varphi(K_n^r) = \varphi(K_n)$. Therefore, by Theorem 1.8, $\varphi(K_n^r) = \frac{1}{2}(n-1)(n-2)$. □

The following results discuss about the sparing numbers of the arbitrary powers of the graph classes which are discussed in Section 2.

Proposition 3.2. *For a positive integer $r > 1$, the sparing number of the r-th power of a complete bipartite graph $K_{m,n}$ is $\frac{1}{2}(m+n-1)(m+n-2)$.*

Proof. Since $K_{m,n}^2 = K_{m+n}$, we have $K_{m,n}^r = K_{m+n}$ for any positive integer $r \geq 2$. Therefore, $\varphi(K_{m,n}^r) = \varphi(K_{m+n}) = \frac{1}{2}(m+n-1)(m+n-2)$. □

Proposition 3.3. *Let G be a split graph, without isolated vertices, that contains a clique K_r and an independent set S with $|S| = s$. Then, for $r \geq 3$, the sparing number of G^r is $\frac{1}{2}(r+s-1)(r+s-2)$.*

Proof. Since S has no isolated vertices in G, every pair vertices of S are at a distance at most 3 among themselves. Hence, G^3 is a complete graph. Therefore, For any $r \geq 3$, G^r is a complete graph. Hence by Theorem 1.8, the sparing number of G^r is $\frac{1}{2}(r+s-1)(r+s-2)$. □

Theorem 3.4. *For a positive integer $r > 2$, the sparing number of H_n^r is*

$$\varphi(H_n^r) = \begin{cases} \lfloor \frac{n}{2} \rfloor (n+3) & \text{if } r = 3 \\ n(2n-1) & \text{if } r \geq 4. \end{cases}$$

Proof. Let u be the central vertex, $V = \{v_1v_2v_3, \ldots, v_n\}$ be the set of vertices of the cycle C_n and $W = \{w_1, w_2, w_3, \ldots,$
$w_n\}$ be the set of pendant vertices in H_n. In H_n, the central vertex u is adjacent to each vertex v_i of V and each v_i is adjacent to a vertex w_i in W.

Since each vertex w_i in W is at a distance at most 3 from u as well as from all vertices of V, for $1 \leq i \leq n$, and from two vertices w_{i-1} and w_{i+1} of W, the subgraph of H_n^3 induced by $V \cup \{u, w_{i-1}, w_i, w_{i+1}\}$ is a complete graph. Hence only one vertex of this set can have a non-singleton set-label. We get minimum number of mono-indexed edges if we label possible number of vertices in W by non-singleton sets. Since w_i is adjacent to w_{i-1} and w_{i+1}, only alternate vertices in W can be labeled by non-singleton sets. Therefore, $\lfloor \frac{n}{2} \rfloor$ vertices in W can be labeled by non-singleton sets. Therefore, since each w_i is of degree $n+3$, total number of edges in H_n^3, that are not mono-indexed, is $\lfloor \frac{n}{2} \rfloor (n+3)$.

The distance between any two points of a helm graph is at most 4. Hence, G^4 is a complete graph. Therefore, For any $r \geq 4$, G^r is a complete graph. Hence by Theorem 1.8, the sparing number of G^r is $n(2n-1)$. □

We have not determined the sparing number of arbitrary powers of paths and cycles yet. The following results discusses the sparing number of the r-th power of a path on n vertices.

The diameter of a path P_m on $n = m + 1$ vertices is $m = n - 1$. Therefore, by Theorem 1.10, $P_m^m = P_{n-1}^{n-1}$ is a complete graph. Hence, we need to study about the r-th powers of P_{n-1} if $r < n - 1$.

Theorem 3.5. *Let P_{n-1} be a path graph on n vertices. Then, its spring number is $\frac{r-1}{2(r+1)}[r(2n-1-r)+2i]$.*

Proof. Let $P_m : v_1v_2v_3 \dots v_n$, where $m = n - 1$. In P_m^2, $d(v_1) = d(v_n) = r, d(v_2) = d(v_{n-1}) = r+1, \dots, d(v_r) = d(v_{n-r+1} = r+r-1 = 2r-1$ and $d(v_j) = 2r,\ r+1 \leq j \leq n-r$. Hence, $\sum_{v \in V(P_n)} d(v) = 2[r+(r+1)+(r+2)+\dots+2r-1)]+(n-2r)2r = r(2n-1-r)$. Therefore, $|E(P_m^r)| = \frac{r}{2}(2n-1-r)$.

It can be seen that among any $r+1$ consecutive vertices $v_i, v_{i+1}, \dots v_{i+r}$ of P_m, r vertices must be mono-indexed. Hence, label $v_1, v_2, \dots, v_k$ by singleton sets and v_{r+1} by a non-singleton set. Since $v_{r+2}, v_{r+3} \dots, v_{2r+1}$ are adjacent to v_{r+1}, they can be labeled only by distinct singleton sets that are not used before for labeling. Now, v_{2r+2} can be labeled by a non-singleton set that has not already been used. Proceeding like this the vertices which has the form $v_{(r+1)k}, (r+1)k \leq n$ can be labeled by distinct non-singleton sets and all other vertices by singleton sets.

If $n \equiv\ i\ (mod\ (k+1))$, then v_{n-i} can also be labeled by a non-singleton set. Then the number of vertices that are not mono-indexed is $\frac{n-i}{r+1}$. Therefore, the number of edges that are not mono-indexed is $2r[\frac{(n-i)}{r+1} - 1] + (r+i) = \frac{1}{r+1}[r(2n-1-r)-(r-1)i]$. Therefore, the total number of mono-indexed edges is $\frac{r}{2}(2n-1-r) - \frac{1}{r+1}[r(2n-1-r)-(r-1)i] = \frac{r-1}{2(r+1)}[r(2n-1-r)+2i]$. □

Figure 5 depicts the cube of a path with a weak IASI defined on it.



Figure 5: Cubes of a path which admits a weak IASI

The diameter of a cycle C_n is $\lfloor \frac{n}{2} \rfloor$. Therefore, by Theorem 1.10, $C_n^{\lfloor \frac{n}{2} \rfloor}$ (and higher powers) is a complete graph. Hence, we need to study about the r-th power of C_n if $r < \lfloor \frac{n}{2} \rfloor$. The following theorem discusses about the sparing number of an arbitrary power of a cycle.

Theorem 3.6. *Let C_n be a cycle on n vertices and let r be a positive integer less than $\lfloor \frac{n}{2} \rfloor$. Then the sparing number of the the r-th power of C_n is given by $\varphi(C_n^r) = \frac{r}{r+1}((r-1)n + 2i)$ if $n \equiv i \ (mod \ (r+1))$.*

Proof. Let $C_n : v_1v_2v_3 \dots v_nv_1$ be the given cycle on n vertices. The graph C_n^r is a $2r$-regular graph. Therefore, we have $|E(C_n^r)| = \frac{1}{2}\sum_{v \in V} d(v) = rn$.

First, label the vertex v_1 in C_n^r by a non-singleton set. Therefore, $2r$ vertices $v_2, v_3, \dots v_{r+1}, v_n, v_{n-1} \dots v_{n-r+1}$ can be labeled only by distinct singleton sets. Next, we can label the vertex v_{r+2} by a non-singleton set, that is not already used for labeling. Since the vertices $v_2, v_3, \dots v_{r+1}$ have already been mono-indexed, r vertices $v_{r+3}, v_{r+4}, \dots v_{2r+2}$ that are adjacent to v_{r+2} in C_n^r must be labeled by distinct singleton sets. Proceeding like this, we can label all the vertices of the form $v_{(r+1)k+1}$, where k is a positive integer less than $\lfloor n \rfloor$, such that $(r+1)k+1 \leq n-r$ (since the last vertex that remains unlabeled is v_{n-r}).

If $n \equiv i \ (mod \ (k+1))$, then $n - i = (r+1)k + 1$ for some positive integer k. Then, $v_{n-(r-i)}$ can be labeled by a non-singleton set. Therefore, the number of vertices that are labeled by non-singleton set is $\frac{n-i}{r+1}$. Since C_n^r is $2r$-regular, the number of edges that are not mono-indexed in C_n^r is $2r\frac{n-i}{r+1}$. Hence, the number of mono-indexed edges is $rn - 2r\frac{n-i}{r+1} = \frac{r}{r+1}((r-1)n + 2i)$. □

Figure 6 illustrates the admissibility of weak IASIs by the squares of even and odd cycles.



Figure 6: Cube of a cycle with a weak IASI defined on it.

4 Conclusion

In this paper, we have established some results on the admissibility of weak IASIs by certain graphs and graph powers. The admissibility of weak IASI by various graph classes, graph operations and graph products and finding the corresponding sparing numbers are still open.

In this paper, we have not addressed the following problems, which are still open. The adjacency and incidence patterns of elements of the graph concerned will matter in determining its admissibility of weak IASI and the sparing number.

Problem 1. Find the sparing number of the r-th power of trees and in particular, binary trees for applicable values of r.

Problem 2. Find the sparing number of the r-th power of bipartite graph and in general, graphs that don't have a complete bipartite graphs as their subgraphs, for applicable values of r.

Problem 3. Find the sparing number of the r-th power of an n-sun graph that is not complete, for applicable values of r.

Problem 4. Find the sparing number of the square of a split graph that is not complete.

Some other standard graph structures related to paths and cycles are lobster graph, ladder graphs, grid graphs and prism graphs. Hence, the following problems are also worth studying.

Problem 5. Find the sparing number of arbitrary powers of a lobster graph.

Problem 6. Find the sparing number of arbitrary powers of a ladder graphs L_n.

Problem 7. Find the sparing number of arbitrary powers of grid graphs (or lattice graphs) $L_{m,n}$.

Problem 8. Find the sparing number of arbitrary powers of prism graphs and anti-prism graphs.

Problem 9. Find the sparing number of arbitrary powers of armed crowns and dragon graphs.

More properties and characteristics of different IASIs, both uniform and non-uniform, are yet to be investigated. The problems of establishing the necessary and sufficient conditions for various graphs and graph classes to have certain IASIs are also open.

References

[1] B. D. Acharya, *Set-Valuations and Their Applications*, MRI Lecture Notes in Applied Mathematics, No. 2, The Mehta Research Institute of Mathematics and Mathematical Physics, Allahabad, 1983.

[2] A. Brandstädt, V. B. Le and J. P. Spinrad, *Graph Classes:A Survey*, SIAM, Philadelphia, 1999.

[3] J. A. Bondy and U. S. R. Murty, *Graph Theory*, Springer, 2008.

[4] J. A. Gallian, *A Dynamic Survey of Graph Labelling*, The Electronic Journal of Combinatorics (DS-6), 2013.

[5] K. A. Germina and T. M. K Anandavally, *Integer Additive Set-Indexers of a Graph:Sum Square Graphs*, Journal of Combinatorics, Information and System Sciences, **37**(2-4)(2012), 345-358.

[6] K. A. Germina and N. K. Sudev, *On Weakly Uniform Integer Additive Set-Indexers of Graphs*, International Mathematical Forum **8**(37)(2013), 1827-34., DOI: 10.12988/imf.2013.310188.

[7] F. Harary, *Graph Theory*, Addison-Wesley Publishing Company Inc., 1994.

[8] N K Sudev and K A Germina, *On Integer Additive Set-Indexers of Graphs*,Int. J. Math. Sci. & Engg. Appl., **8**(2)(2014), 11-22.

[9] N. K. Sudev and K. A. Germina, *A Characterisation of Weak Integer Additive Set-Indexers of Graphs*,Journal of Fuzzy Set Valued Analysis, **2014**(2014), Article Id:jfsva-00189, 1-7., DOI: 10.5899/2014/jfsva-00189.

[10] N. K. Sudev and K. A. Germina, *Weak Integer Additive Set-Indexers of Certain Graph Operations*, Global Journal of Mathematical Sciences: Theory and Practical, **6**(2)(2014), 25-36.

[11] N. K. Sudev and K. A. Germina, *A Note on the Sparing Number of Graphs*, Advances and Applications in Discrete Mathematics, **14**(1)(2014), 51-65.

[12] N. K. Sudev and K. A. Germina, *On the Sparing Number of Certain Graph Classes*, Journal of Discrete Mathematical Sciences nd Cryptography, **18**(1-2)(2015), 117-128, DOI: 10.1080/09720529.2014.962866.

[13] N. K. Sudev and K. A. Germina, *On the Sparing Number of Certain Graph Structures*, Annals of Pure and Applied Mathematics, **6**(2)(2014), 140-149.

[14] E. W. Weisstein, *CRC Concise Encyclopedia of Mathematics*, CRC press, 2011.

[15] D. B. West, *Introduction to Graph Theory*, Pearson Education Inc., 2001.

[16] Information System on Graph Classes and their Inclusions `www.graphclasses.org`